

Native Porcine α-Ketoglutarate Dehydrogenase

Cat. No. NATE-0495

Lot. No. (See product label)

Introduction

Description The oxoglutarate dehydrogenase complex (OGDC) or α -ketoglutarate dehydrogenase complex is an

enzyme complex, most commonly known for its role in the citric acid cycle.

Synonyms 2-ketoglutarate dehydrogenase; 2-oxoglutarate dehydrogenase; 2-oxoglutarate:lipoate oxidoreductase; 2-

oxoglutarate:lipoamide 2-oxidoreductase (decarboxylating and acceptor-succinylating); α -ketoglutarate dehydrogenase; α -ketoglutaric dehydrogenase; α -oxoglutarate dehydrogenase; AKGDH; OGDC; ketoglutaric dehydrogenase; oxoglutarate decarboxylase; oxoglutarate

dehydrogenase; oxoglutarate dehydrogenase (lipoamide); EC 1.2.4.2; 9031-02-1

Product Information

Species Porcine

Source Porcine heart

Form buffered aqueous glycerol solution. Supplied as a 50% glycerol solution containing ~9 mg per mL bovine

serum albumin, 30% sucrose, 1.5 mM EDTA, 1.5 mM EGTA, 1.5 mM 2-mercaptoethanol, 0.3% TRITON™ X-

100, 0.003% sodium azide, and 15 mM potassium phosphate, pH 6.8.

EC Number EC 1.2.4.2

CAS No. 9031-02-1

Activity 0.1-1.0 units/mg protein (Lowry)

Unit One unit will convert 1.0 μmole of β -NAD to β -NADH per min at pH 7.4 at 30°C in the presence of

Definition saturating levels of coenzyme A.

Storage and Shipping Information

Storage −20°C

Tel: 1-631-562-8517 1-516-512-3133 **Email:** info@creative-enzymes.com

1/1