

Native Human Superoxide Dismutase

Cat. No. NATE-0680

Lot. No. (See product label)

Introduction

Description Superoxide dismutase (SOD) catalyzes the dismutation of superoxide radicals to hydrogen peroxide and molecular oxygen. SOD plays a critical role in the defense of cells against the toxic effects of oxygen radicals. SOD competes with nitric oxide (NO) for superoxide anion (which reacts with NO to form peroxynitrite), thereby SOD promotes the activity of NO. SOD has also been shown to suppress apoptosis in cultured rat ovarian follicles, neural cell lines, and transgenic mice by preventing the conversion of NO to peroxynitrate, an inducer of apoptosis.

Applications Superoxide dismutase from human erythrocytes has been used in a study to identify in vitro glycosylated sites of human Cu-Zn-superoxide dismutase. Superoxide dismutase from human erythrocytes has also been used in a study to investigate a prospective test-system for the screening of cytoprotective drugs and their combinations.

Synonyms Superoxide dismutases; EC 1.15.1.1; superoxidase dismutase; copper-zinc superoxide dismutase; Cu-Zn superoxide dismutase; ferrisuperoxide dismutase; superoxide dismutase I; superoxide dismutase II; SOD; Cu,Zn-SOD; Mn-SOD; Fe-SOD; SODF; SODS; SOD-1; SOD-2; SOD-3; SOD-4; hemocuprein; erythrocuprein; cytocuprein; cuprein ; hepatocuprein; 9054-89-1

Product Information

Species Human

Source Human erythrocytes

Form Lyophilized powder containing potassium phosphate buffer salts

EC Number EC 1.15.1.1

CAS No. 9054-89-1

Activity > 2,500 units/mg protein

Pathway Amyotrophic lateral sclerosis (ALS), organism-specific biosystem; Amyotrophic lateral sclerosis (ALS), conserved biosystem; FOXA1 transcription factor network, organism-specific biosystem; Folate Metabolism, organism-specific biosystem; Hemostasis, organism-specific biosystem; Huntingtons disease, organism-specific biosystem; Huntingtons disease, conserved biosystem; FoxO family signaling, organism-specific biosystem; Huntingtons disease, organism-specific biosystem; Huntingtons disease, conserved biosystem; Oxidative Stress, organism-specific biosystem; Peroxisome, organism-specific biosystem; Peroxisome, conserved biosystem; Selenium Pathway, organism-specific biosystem; Folate Metabolism, organism-specific biosystem; Oxidative Stress, organism-specific biosystem; Selenium Pathway, organism-specific biosystem; superoxide radicals degradation, organism-specific biosystem

Function chaperone binding; copper ion binding; metal ion binding; oxidoreductase activity; protein binding; protein homodimerization activity; protein phosphatase 2B binding; superoxide dismutase activity; zinc ion binding; DNA binding; identical protein binding; manganese ion binding; manganese ion binding; metal ion binding; oxidoreductase activity; oxygen binding; superoxide dismutase activity; superoxide dismutase activity; copper ion binding; heparin binding; metal ion binding; oxidoreductase activity; protein binding; superoxide dismutase activity; zinc ion binding

Unit One unit will inhibit reduction of Cytochrome c by 50% in a coupled system with xanthine oxidase at pH

Unit

One unit will inhibit reduction of cytochrome c by 50% in a coupled system with xanthine oxidase at pH 7.8 at 25°C in a 3.0 mL reaction volume. Xanthine oxidase concentration should produce an initial ΔA_{550} of 0.025 ± 0.005 per min.

Storage and Shipping Information

Storage -20°C