

β-N-Acetylglucosaminidase from *Streptococcus pneumoniae*, Recombinant

Cat. No. NATE-0782

Lot. No. (See product label)

Introduction

Description Hexosaminidase, sometimes called β-N-acetylhexosaminidase, is reported to liberate terminal β-linked N-acetylglucosamine and N-acetylgalactosamine from a variety of substrates. The activity of β-N-acetylglucosaminidase may be determined with the chromogenic substrate p-nitrophenyl-N-acetyl-β-D-glucosaminide. β-N-acetylglucosaminidase hydrolyzes the terminal nonreducing N-acetyl-D-hexosamine residues. This enzyme contains two predominant isozymes, Hex A, a heterodimer, and Hex B, a homodimer. N-acetylglucosamine, acetamide, N-2-acetamido-2-deoxyglucosylamine, N-acetylnojirimycin, and N-acetyldeoxynojirimycin are known inhibitors.

Applications Hexosaminidase is an enzyme involved in the hydrolysis of terminal N-acetyl-D-hexosamine residues in N-acetyl-β-D-hexosaminides.

Synonyms EC 3.2.1.52; 9012-33-3; hexosaminidase; β-acetylaminodeoxyhexosidase; N-acetyl-β-D-hexosaminidase; N-acetyl-beta-hexosaminidase; β-hexosaminidase; β-acetylhexosaminidase; β-D-N-acetylhexosaminidase; β-N-acetyl-D-hexosaminidase; β-N-acetylglucosaminidase; hexosaminidase A; N-acetylhexosaminidase; β-D-hexosaminidase

Product Information

Species *Streptococcus pneumoniae*

Source *E. coli*

Form buffered aqueous solution, Solution in 20 mM Tris-HCl, pH 7.5, 25 mM NaCl

EC Number 3.2.1.52

CAS No. 9012-33-3

Activity > 80 units/mg protein

Unit Definition One unit will hydrolyze 1.0 μmole of p-nitrophenyl N-acetyl-β-D-glucosaminide to p-nitrophenol and N-acetyl-D-glucosamine per min at the pH 5 at 37°C.

Usage and Packaging

Package vial of > 1.0 unit

Storage and Shipping Information

Storage 2-8°C