

## Transglutaminase 2 from Rabbit, Recombinant

Cat. No. NATE-1734

Lot. No. (See product label)

### Introduction

<b>Description</b>	This enzyme is based on the NCBI database sequence XM_008256006. It is N-terminally fused to a hexahistidine-tag resulting in the encoded N-terminal amino acid sequence MHHHHHHAEDLIL.... This enzyme is produced in E. coli and purified by ion metal chelating chromatography to more than 95 % purity.
<b>Applications</b>	The transglutaminase 2 catalyzes acyl transfer reactions from glutamin residues in proteins or peptides to primary amines, e. g. the formation of $\epsilon$ -( $\gamma$ -glutamyl) lysine bonds between proteins by transferring the acyl group of a peptide-bound glutamine residue to the primary amino group of a peptide-bound lysine residue. The transglutaminase 2 may also be used for immunoprecipitation.
<b>Synonyms</b>	transglutaminase; EC 2.3.2.13; 80146-85-6; transglutaminase; Factor XIIIa; fibrinolygase; fibrin stabilizing factor; glutaminylpeptide $\gamma$ -glutamyltransferase; polyamine transglutaminase; tissue transglutaminase; R-glutaminy-peptide:amine $\gamma$ -glutamyl transferase; protein-glutamine $\gamma$ -glutamyltransferase; TG1

### Product Information

<b>Species</b>	Rabbit
<b>Source</b>	E. coli
<b>Appearance</b>	White lyophilized solid.
<b>Form</b>	The enzyme is lyophilized from 50 mM NaH <sub>2</sub> PO <sub>4</sub> , 150 mM NaCl, pH 8.0. Sample contains maltodextrin.
<b>EC Number</b>	EC 2.3.2.13
<b>CAS No.</b>	80146-85-6
<b>Molecular Weight</b>	77 kDa
<b>Purity</b>	> 95 % (visually by SDS-PAGE)
<b>Activity</b>	> 750 U/mg [Activity is determined by measuring the rate of fluorescence enhancement after transglutaminase-catalyzed monodansylcadaverine-incorporation into N,N-dimethylated casein according to Lorand et al., Anal. Biochem. 44 (221-231)].
<b>Activators</b>	Add 10 mM Ca <sup>2+</sup> to activate transglutaminase.
<b>Unit Definition</b>	1 U is defined as the increase in fluorescence intensity of 1 a.u./min (measured on a Cary eclipse fluorescence spectrophotometer, Varian; $\lambda_{ex}$ = 332 nm, $\lambda_{em}$ = 500 nm; band filter = 10 nm; detector strength = 600 V; temperature = 37 °C, assay volume = 1 ml)].

### Usage and Packaging

<b>Package</b>	250 $\mu$ g; 1mg
<b>Reconstitution</b>	Add the volume of H <sub>2</sub> O the protein is lyophilized from to the vial of lyophilized powder. Rotate vial gently until solid dissolves. After reconstitution, the solution should be stored frozen in working aliquots. Keep cooled on ice for short term storage.

### ***Storage and Shipping Information***

#### ***Storage***

Store at -20 °C in working aliquots. Repeated freezing and thawing is not recommended. Delivery is possible at ambient temperature.