

Native Sphingomyelinase from Streptomyces sp.

Cat. No. NATE-1160

Lot. No. (See product label)

Introduction

| Description | Sphingomyelin phosphodiesterase is a hydrolase enzyme that is involved in sphingolipid metabolism reactions. SMase is a member of the DNase I superfamily of enzymes and is responsible for breaking sphingomyelin (SM) down into phosphocholine and ceramide. The activation of SMase has been suggested as a major route for the production of ceramide in response to cellular stresses. |
|--------------|---|
| Applications | This enzyme is useful for enzymatic determination of sphingomyelin when coupled With alkaline phosphatase and choline oxidase. |
| Synonyms | Sphingomyelin phosphodiesterase; EC 3.1.4.12; neutral sphingomyelinase; 9031-54-3; sphingomyelin cholinephosphohydrolase; sphingomyelinase; Smase |

Product Information

| Source | Streptomyces sp. |
|-----------------------|--|
| Appearance | White to brownish amorphous powder, lyophilized |
| Form | Freeze dried powder |
| EC Number | EC 3.1.4.12 |
| CAS No. | 9031-54-3 |
| Molecular Weight | 37.5 kDa (SDS–PAGE) |
| Activity | > 500 U/mg |
| lsoelectric point | 8.6 |
| pH Stability | 5.0-8.0 |
| Optimum pH | 7.0–8.0 |
| Thermal stability | Stable at 40°C and below |
| Michaelis Constant | Sphingomyelin $0.45 \times 10-3M$ |
| Activators | Mg2+, Mn2+, Non–ionic detergents |
| Inhibitors | EDTA |
| Stabilizers | Mg2+ |
| Unit Definition | One unit is defined as the amount of enzyme which hydrolyzes 1 μ mole of sphingomyelin per minute at 37°C under the conditions specified in the assay procedure. |

Storage and Shipping Information

Storage Storage at -20°C in the presence of a desiccant is recommended.

Stability At least one year at -20°C