

Native Human Plasmin

Cat. No. NATE-0598 Lot. No. (See product label)

Introduction	
Description	Plasmin functions as a key enzyme of the fibrinolytic cascade, and is also important in inflammation processes. Plasmin exhibits preferential cleavage at the carboxyl side of lysine and arginine residues with higher selectivity than trypsin. Converts polymerized fibrin into soluble products.
Applications	A complex between plasmin and an inhibitor has been isolated in a study via affinity chromatography from urokinase-activated human plasma. It has also been used in a study to investigate activation of human epithelial sodium channel (ENaC) by plasmin and chymotrypsin.
Synonyms	fibrinase; fibrinolysin; actase; serum tryptase; thrombolysin; plasmin; EC 3.4.21.7; 9001-90-5; PLG
Product Information	
Species	Human
Source	Human plasma
Form	Lyophilized powder containing sodium phosphate, mannitol, and NaCl.
EC Number	EC 3.4.21.7
CAS No.	9001-90-5
Activity	> 2.0 units/mg protein
Optimum pH	8.5
Optimum temperature	37°C with rapid inactivation at 56°C
Pathway	Activation of Matrix Metalloproteinases, organism-specific biosystem; Angiopoietin receptor Tie2-mediated signaling, organism-specific biosystem; Blood Clotting Cascade, organism-specific biosystem; Complement and Coagulation Cascades, organism-specific biosystem; Complement and coagulation cascades, organism- specific biosystem; Complement and coagulation cascades, organism- specific biosystem; Complement and coagulation cascades, conserved biosystem; Degradation of the extracellular matrix, organism-specific biosystem
Function	apolipoprotein binding; cell surface binding; peptidase activity; protein binding; protein domain specific binding; serine-type endopeptidase activity
Unit Definition	One unit will produce one $\mu mole$ of p-Nitroanilide from D-Val-Leu-Lys-p-Nitroanilide per minute at pH 7.5 at 37°C.
Storage and Shipping Information	

Storage

-20°C