

Native *Geobacillus stearothermophilus* Thermolysin

Cat. No. NATE-0704

Lot. No. (See product label)

Introduction

Description

Thermolysin is a thermostable neutral metalloproteinase enzyme produced by the Gram-positive bacteria *Bacillus thermoproteolyticus*. It requires one zinc ion for enzyme activity and four calcium ions for structural stability. Thermolysin specifically catalyzes the hydrolysis of peptide bonds containing hydrophobic amino acids. However thermolysin is also widely used for peptide bond formation through the reverse reaction of hydrolysis. Thermolysin is the most stable member of a family of metalloproteinases produced by various *Bacillus* species. These enzymes are also termed 'neutral' proteinases or thermolysin-like proteinases (TLPs).

Applications

A thermostable (thermophilic) extracellular metalloendopeptidase containing four calcium ions. Cofactors are zinc and calcium. Hydrolyzes protein bonds on the N-terminal side of hydrophobic amino acid residues. The pH optimum is 8.0 and the optimal temperature for activity is 70°C. Considerably stable from pH 5 to 9.5. Thermolysin has a low cleavage specificity, therefore, it produces a number of short fragments that are suitable for sequencing. Preferential cleavage: X-cleavage-Y-Z where X=any amino acid; Y=Leu, Phe, Ile, Val, Met, Ala and Z is any amino acid other than Pro. Cleavage N-terminal to Leu is preferred over cleavage of N-terminal to Phe which is preferred over the others. Often used to do limited proteolysis for peptide mapping and studies of protein structure and conformational changes.

Synonyms

thermolysin; *Bacillus thermoproteolyticus* neutral proteinase; thermoase; thermoase Y10; TLN; EC 3.4.24.27

Product Information

Source

Geobacillus stearothermophilus

Form

lyophilized powder containing calcium and sodium acetate buffer salts

EC Number

EC 3.4.24.27

CAS No.

9073-78-3

Activity

30-175 units/mg protein (E1%/280)

Unit Definition

One unit will hydrolyze casein to produce color equivalent to 1.0 μ mole (181 μ g) of tyrosine per min at pH 7.5 at 37°C.

Storage and Shipping Information

Storage

-20°C