

D-arabinitol dehydrogenase (NADP+)

Cat. No. EXWM-0195

Lot. No. (See product label)

Introduction

Description The enzyme from the rust fungus Uromyces fabae can use D-arabinitol and D-mannitol as substrates in the forward direction and D-xylulose, D-ribulose and, to a lesser extent, D-fructose as substrates in the reverse direction. This enzyme carries out the reactions of both EC 1.1.1.11, D-arabinitol 4-dehydrogenase and EC 1.1.1.250, D-arabinitol 2-dehydrogenase, but unlike them, uses NADP+ rather than NAD+ as cofactor. D-Arabinitol is capable of quenching reactive oxygen species involved in defense reactions of the host plant.

Synonyms NADP+-dependent D-arabitol dehydrogenase; ARD1p; D-arabitol dehydrogenase 1

Product Information

Form	Liquid or lyophilized powder
EC Number	EC 1.1.1.287
Reaction	(1) D-arabinitol + NADP+ = D-xylulose + NADPH + H+; (2) D-arabinitol + NADP+ = D-ribulose + NADPH + H+
Notes	This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.
Storage and Shipping Information	

Storage Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.