

2-hydroxy-4-carboxymuconate semialdehyde hemiacetal dehydrogenase

Cat. No. EXWM-0224

Lot. No. (See product label)

Introduction

Description

The enzyme does not act on unsubstituted aliphatic or aromatic aldehydes or glucose; NAD⁺ can replace NADP⁺, but with lower affinity. The enzyme was initially believed to act on 4-carboxy-2-hydroxy-cis,cis-muconate 6-semialdehyde and produce 4-carboxy-2-hydroxy-cis,cis-muconate. However, later studies showed that the substrate is the hemiacetal form, and the product is 2-oxo-2H-pyran-4,6-dicarboxylate.

Synonyms

2-hydroxy-4-carboxymuconate 6-semialdehyde dehydrogenase; 4-carboxy-2-hydroxy-cis,cis-muconate-6-semialdehyde:NADP⁺ oxidoreductase; α-hydroxy-γ-carboxymuconic ε-semialdehyde dehydrogenase; 4-carboxy-2-hydroxymuconate-6-semialdehyde dehydrogenase; LigC; ProD

Product Information

Form

Liquid or lyophilized powder

EC Number

EC 1.1.1.312

Reaction

4-carboxy-2-hydroxymuconate semialdehyde hemiacetal + NADP⁺ = 2-oxo-2H-pyran-4,6-dicarboxylate + NADPH + H⁺

Notes

This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage

Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.