

## monocyclic monoterpene ketone monooxygenase

Cat. No. EXWM-0704

Lot. No. (See product label)

### Introduction

#### Description

A flavoprotein (FAD). This Baeyer-Villiger monooxygenase enzyme from the Gram-positive bacterium *Rhodococcus erythropolis* DCL14 has wide substrate specificity, catalysing the lactonization of a large number of monocyclic monoterpene ketones and substituted cyclohexanones. Both (1R,4S)- and (1S,4R)-1-hydroxymenth-8-en-2-one are metabolized, with the lactone product spontaneously rearranging to form 3-isopropenyl-6-oxoheptanoate.

#### Synonyms

1-hydroxy-2-oxolimonene 1,2-monooxygenase; dihydrocarvone 1,2-monooxygenase; MMKMO

### Product Information

#### Form

Liquid or lyophilized powder

#### EC Number

EC 1.14.13.105

#### Reaction

(1) (-)-menthone + NADPH + H<sup>+</sup> + O<sub>2</sub> = (4R,7S)-7-isopropyl-4-methyloxepan-2-one + NADP<sup>+</sup> + H<sub>2</sub>O; (2) dihydrocarvone + NADPH + H<sup>+</sup> + O<sub>2</sub> = 4-isopropenyl-7-methyloxepan-2-one + NADP<sup>+</sup> + H<sub>2</sub>O; (3) (iso)-dihydrocarvone + NADPH + H<sup>+</sup> + O<sub>2</sub> = 6-isopropenyl-3-methyloxepan-2-one + NADP<sup>+</sup> + H<sub>2</sub>O; (4a) 1-hydroxymenth-8-en-2-one + NADPH + H<sup>+</sup> + O<sub>2</sub> = 7-hydroxy-4-isopropenyl-7-methyloxepan-2-one + NADP<sup>+</sup> + H<sub>2</sub>O; (4b) 7-hydroxy-4-isopropenyl-7-methyloxepan-2-one = 3-isopropenyl-6-oxoheptanoate (spontaneous)

#### Notes

This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

### Storage and Shipping Information

#### Storage

Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.