

nitric-oxide synthase [NAD(P)H]

Cat. No. EXWM-0764

Lot. No. (See product label)

Introduction

Description Binds heme (iron protoporphyrin IX) and tetrahydrobiopterin. Most of the bacterial and archaeal enzymes consist of only an oxidase domain and function together with bacterial ferredoxins. The enzyme from the Δ -proteobacterium *Sorangium cellulosum* also includes a reductase domain that binds FAD, FMN and a [2Fe-2S] cluster. The similar enzymes from plants and animals use only NADPH as acceptor (cf. EC 1.14.13.39).

Synonyms nitric oxide synthetase; NO synthase

Product Information

Form Liquid or lyophilized powder

EC Number EC 1.14.13.165

Reaction $2 \text{ L-arginine} + 3 \text{ NAD(P)H} + 3 \text{ H}^+ + 4 \text{ O}_2 = 2 \text{ L-citrulline} + 2 \text{ nitric oxide} + 3 \text{ NAD(P)}^+ + 4 \text{ H}_2\text{O}$ (overall reaction); (1a) $2 \text{ L-arginine} + 2 \text{ NAD(P)H} + 2 \text{ H}^+ + 2 \text{ O}_2 = 2 \text{ N}\omega\text{-hydroxy-L-arginine} + 2 \text{ NAD(P)}^+ + 2 \text{ H}_2\text{O}$; (1b) $2 \text{ N}\omega\text{-hydroxy-L-arginine} + \text{NAD(P)H} + \text{H}^+ + 2 \text{ O}_2 = 2 \text{ L-citrulline} + 2 \text{ nitric oxide} + \text{NAD(P)}^+ + 2 \text{ H}_2\text{O}$

Notes This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.