

acyl-lipid ω -6 desaturase (cytochrome b5)

Cat. No. EXWM-0985

Lot. No. (See product label)

Introduction

Description

This microsomal enzyme introduces a cis double bond in fatty acids attached to lipid molecules at a location 6 carbons away from the methyl end of the fatty acid. The distance from the carboxylic acid end of the molecule does not affect the location of the new double bond. The most common substrates are oleoyl groups attached to either the sn-1 or sn-2 position of the glycerol backbone in phosphatidylcholine. cf. EC 1.14.19.23, acyl-lipid ω -6 desaturase (ferredoxin).

Synonyms

oleate desaturase (ambiguous); linoleate synthase (ambiguous); oleoyl-CoA desaturase (incorrect); oleoylphosphatidylcholine desaturase (ambiguous); phosphatidylcholine desaturase (ambiguous); n-6 desaturase (ambiguous); FAD2 (gene name)

Product Information

Form

Liquid or lyophilized powder

EC Number

EC 1.14.19.22

CAS No.

72536-70-0

Reaction

an oleoyl-[glycerolipid] + 2 ferrocytochrome b5 + O₂ + 2 H⁺ = a linoleoyl-[glycerolipid] + 2 ferricytochrome b5 + 2 H₂O

Notes

This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage

Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.