

## Native Penicillium sp. Dextranase

Cat. No. NATE-0194

Lot. No. (See product label)

### Introduction

**Description** An endodextranase that hydrolyzes-(1,6)-glucosidic linkages in dextran. Dextranase is an undesirable compound synthesized from sucrose by microbial contaminants during sugar production that increases viscosity of the flow and decreases industrial recovery. Dextranase has been used for hydrolyzing dextran at sugar mills in order to improve efficiency of sugar production.

**Applications** Dextranase from *Penicillium* has been used in a study to assess the purification properties of an extracellular dextranase from *Penicillium janthinellum*. Dextranase from *Penicillium* has also been used in a study to investigate the carbohydrate component of *Penicillium funiculosum* dextranase. It has been used for the hydrolysis of carbohydrate polymers, during the study of polysaccharide synthesis by *Phanerochaete chrysosporium*. It has also been used in the synthesis of new enzymatically degradable thermo-responsive nanogels.

**Synonyms** EC 3.2.1.11, dextran hydrolase; endodextranase; dextranase DL 2; DL 2; endo-dextranase;  $\alpha$ -D-1,6-glucan-6-glucanohydrolase; 1,6- $\alpha$ -D-glucan 6-glucanohydrolase; 9025-70-1; Dextranase

### Product Information

<b>Source</b>	Penicillium sp.
<b>Form</b>	lyophilized powder.
<b>EC Number</b>	EC 3.2.1.11
<b>CAS No.</b>	9025-70-1
<b>Activity</b>	3,000 units/mg
<b>Optimum pH</b>	5
<b>Optimum temperature</b>	50°C

### Storage and Shipping Information

<b>Storage</b>	2-8°C
----------------	-------