

Native Porcine Diamine Oxidase

Cat. No. NATE-0189 Lot. No. (See product label)

Introduction

Description	Diamine oxidase from porcine kidney is a homodimer consisting of 2 equal subunits with a molecular weight of 87 kDa each. Each subunit contains one molecule of pyridoxal phosphate and one atom of copper. The molecular mass of the enzyme is found to be 170 kDa. The enzyme is a glycoprotein
	containing 5% hexose, 3.3% glucosamine, 2.6% N-acetylglucosamine, and 0.25% N-acetylneuraminic acid. The enzyme exhibits a high affinity for concanavalin A. It catalyzes the oxidation of monoamines,
	diamines, and histamine to aldehydes, ammonia, and hydrogen peroxide. Optimum pH with cadverine and histamine as substrates is found to be 6.3-7.4.2 The enzyme is classified as a copper amine oxidase and it is a key enzyme in nitrogen metabolism. It is inhibited by diethyldithiocarbamate,
	phenylhydrazine, semicarbazide, cyanide, isonicotinic acid hydrazide.

- **Applications**An endodextranase that hydrolyzes α -(1,6)-glucosidic linkages in dextran. Dextrans are undesirable
compounds synthesized from sucrose by microbial contaminants during sugar production that increase
viscosity of the flow and decrease industrial recovery. Dextranase has been used for hydrolyzing dextran
at sugar mills in order to improve efficiency of sugar production. Diamine oxidase from porcine kidney
has been used in a study to investigate a luminescence-based test for determining ornithine
decarboxylase activity. Diamine oxidase from porcine kidney has also been used in a study to investigate
a number of sugar production. Number of the study to investigate
a number of the study to investigate a luminescence-based test for determining ornithine
decarboxylase activity. Diamine oxidase from porcine kidney has also been used in a study to investigate
a number oxidase.
- *Synonyms* EC 1.4.3.6; 9001-53-0; Amine:oxygen oxidoreductase (deaminating) (pyridoxal-containing); Diamine Oxidase; Amine oxidase (copper-containing)

Product Information	
Species	Porcine
Source	Porcine kidney
EC Number	EC 1.4.3.6
CAS No.	9001-53-0
Activity	> 0.05 unit/mg solid
Buffer	100 mM sodium phosphate buffer, pH 7.2: soluble 10 mg/mL
Unit Definition	One unit will oxidize 1.0 $\mu mole$ of putrescine per hr at pH 7.2 at 37°C.

Storage and Shipping Information

Storage –20°C