

precorrin-2 dehydrogenase

Cat. No. EXWM-1345

Lot. No. (See product label)

Introduction

Description This enzyme catalyses the second of three steps leading to the formation of siroheme from uroporphyrinogen III. The first step involves the donation of two S-adenosyl-L-methionine-derived methyl groups to carbons 2 and 7 of uroporphyrinogen III to form precorrin-2 (EC 2.1.1.107, uroporphyrin-III C-methyltransferase) and the third step involves the chelation of ferrous iron to sirohydrochlorin to form siroheme (EC 4.99.1.4, sirohydrochlorin ferrochelatase). In *Saccharomyces cerevisiae*, the last two steps are carried out by a single bifunctional enzyme, Met8p. In some bacteria, steps 1-3 are catalysed by a single multifunctional protein called CysG, whereas in *Bacillus megaterium*, three separate enzymes carry out each of the steps, with SirC being responsible for the above reaction.

Synonyms Met8p; SirC; CysG

Product Information

Form Liquid or lyophilized powder

EC Number EC 1.3.1.76

CAS No. 227184-47-6

Reaction precorrin-2 + NAD⁺ = sirohydrochlorin + NADH + H⁺

Notes This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.