

pteridine reductase

Cat. No. EXWM-1515

Lot. No. (See product label)

Introduction

Description The enzyme from Leishmania (both amastigote and promastigote forms) catalyses

the reduction by NADPH of folate and a wide variety of unconjugated pterins, including biopterin, to their tetrahydro forms. It also catalyses the reduction of 7,8-dihydropterins and 7,8-dihydrofolate to their tetrahydro forms. In contrast to EC 1.5.1.3 (dihydrofolate reductase) and EC 1.5.1.34 (6,7-dihydropteridine reductase), pteridine reductase will not catalyse the reduction of the quinonoid form of dihydrobiopterin. The enzyme is specific for NADPH; no activity has been detected with NADH. It also differs from EC 1.5.1.3 (dihydrofolate reductase) in being specific

1/1

for the Si-face of NADPH.

Synonyms PTR1; pteridine reductase 1

Product Information

Form Liquid or lyophilized powder

EC Number EC 1.5.1.33

CAS No. 131384-61-7

Reaction 5,6,7,8-tetrahydrobiopterin + 2 NADP+ = biopterin + 2 NADPH + 2 H+

Notes This item requires custom production and lead time is between 5-9 weeks. We can

custom produce according to your specifications.

Storage and Shipping Information

Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.

Tel: 1-631-562-8517 1-516-512-3133 **Email:** info@creative-enzymes.com