

NADPH-hemoprotein reductase

Cat. No. EXWM-1580

Lot. No. (See product label)

Introduction

Description A flavoprotein containing both FMN and FAD. This enzyme catalyses the transfer of

electrons from NADPH, an obligatory two-electron donor, to microsomal P-450 monooxygenases (e.g. EC 1.14.14.1, unspecific monooxygenase) by stabilizing the one-electron reduced form of the flavin cofactors FAD and FMN. It also reduces cytochrome b5 and cytochrome c. The number n in the equation is 1 if the

hemoprotein undergoes a 2-electron reduction, and is 2 if it undergoes a 1-electron

reduction.

Synonyms CPR; FAD-cytochrome c reductase; NADP-cytochrome c reductase; NADP-

cytochrome reductase; NADPH-dependent cytochrome c reductase; NADPH:P-450 reductase; NADPH:ferrihemoprotein oxidoreductase; NADPH-cytochrome P-450 oxidoreductase; NADPH-cytochrome c oxidoreductase; NADPH-cytochrome c reductase; NADPH-cytochrome p-450 reductase; NADPH-ferricytochrome c oxidoreductase; NADPH-ferrihemoprotein reductase; TPNH2 cytochrome c

reductase; TPNH-cytochrome c reductase; aldehyde reductase (NADPH-dependent);

cytochrome P-450 reductase; cytochrome c reductase (reduced nicotinamide

adenine dinucleotide phosphate, NADPH, NADPH-dependent);

dihydroxynicotinamide adenine dinucleotide phosphate-cytochrome c reductase; ferrihemoprotein P-450 reductase; reduced nicotinamide adenine dinucleotide

phosphate-cytochrome c reductase; reductase, cytochrome c (reduced

nicotinamide adenine dinucleotide phosphate)

Product Information

Form Liquid or lyophilized powder

EC Number EC 1.6.2.4

CAS No. 9023-03-4

Reaction NADPH + H+ + n oxidized hemoprotein = NADP+ + n reduced hemoprotein

Notes This item requires custom production and lead time is between 5-9 weeks. We can

custom produce according to your specifications.

Storage and Shipping Information

Storage Store it at +4 °C for short term. For long term storage, store it at -20 °C \sim -80 °C.

Tel: 1-631-562-8517 1-516-512-3133 **Email:** info@creative-enzymes.com 1/1