

## Native *Candida boidinii* Formate Dehydrogenase

Cat. No. NATE-0254

Lot. No. (See product label)

### Introduction

#### Description

Formate dehydrogenase is involved in the stress response of plants and catalyzes the reduction of NAD<sup>+</sup> to NADH.

#### Applications

Formate Dehydrogenase (FDH) is used for diagnostics in large scale industrial processes. Its used in the production of an unnatural amino acid, tert-L-leucine, a component of some HIV protease and matrix metalloprotease inhibitors.

#### Synonyms

EC 1.2.1.2; 9028-85-7; formate-NAD oxidoreductase; FDH; FDH I; FDH II; N-FDH; formic hydrogen-lyase; formate hydrogenlyase; hydrogenlyase; NAD-linked formate dehydrogenase; NAD-dependent formate dehydrogenase; formate dehydrogenase (NAD); NAD-formate dehydrogenase; formate benzyl-viologen oxidoreductase; formic acid dehydrogenase

### Product Information

#### Source

*Candida boidinii*

#### Form

Type I, lyophilized powder; Type II, powder; Type III, clear brown liquid.

#### EC Number

EC 1.2.1.2

#### CAS No.

9028-85-7

#### Activity

Type I, 5.0-15.0 units/mg protein; Type II, 0.3-0.6 units/mg; Type III, ~50 U/mL.

#### Unit Definition

One unit will oxidize 1.0  $\mu$ mole of formate to CO<sub>2</sub> per min in the presence of  $\beta$ -NAD at pH 7.6 at 37°C.

### Storage and Shipping Information

#### Storage

-20°C