Creative Enzymes®

NAD(P)H dehydrogenase (quinone)

Cat. No. EXWM-1591

Lot. No. (See product label)

Introduction

Description	A flavoprotein. The enzyme catalyses a two-electron reduction and has a preference for short-chain acceptor quinones, such as ubiquinone, benzoquinone, juglone and duroquinone. The animal, but not the plant, form of the enzyme is inhibited by dicoumarol.
Synonyms	menadione reductase; phylloquinone reductase; quinone reductase; dehydrogenase, reduced nicotinamide adenine dinucleotide (phosphate, quinone); DT-diaphorase; flavoprotein NAD(P)H-quinone reductase; menadione oxidoreductase; NAD(P)H dehydrogenase; NAD(P)H menadione reductase; NAD(P)H-quinone dehydrogenase; NAD(P)H-quinone oxidoreductase; NAD(P)H: (quinone-acceptor)oxidoreductase; NAD(P)H: menadione oxidoreductase; NADH-menadione reductase; naphthoquinone reductase; p-benzoquinone reductase; reduced NAD(P)H dehydrogenase; viologen accepting pyridine nucleotide oxidoreductase; vitamin K reductase; diaphorase; reduced nicotinamide-adenine dinucleotide (phosphate) dehydrogenase; vitamin-K reductase; NAD(P)H2 dehydrogenase (quinone); NQO1; QR1; NAD(P)H:(quinone-acceptor) oxidoreductase

Product Information

Form	Liquid or lyophilized powder
EC Number	EC 1.6.5.2
CAS No.	9032-20-6
Reaction	NAD(P)H + H+ + a quinone = NAD(P)+ + a hydroquinone
Notes	This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage	Store it at +4 ℃ for short term. For long term storage, store it at -20 ℃~-80 ℃.