

## nitrite reductase (NO-forming)

Cat. No. EXWM-1616 Lot. No. (See product label)

Introduction	
Description	The reaction is catalysed by two types of enzymes, found in the perimplasm of denitrifying bacteria. One type comprises proteins containing multiple copper centres, the other a heme protein, cytochrome cd1. Acceptors include c-type cytochromes such as cytochrome c-550 or cytochrome c-551 from Paracoccus denitrificans or Pseudomonas aeruginosa, and small blue copper proteins such as azurin and pseudoazurin. Cytochrome cd1 also has oxidase and hydroxylamine reductase activities. May also catalyse the reaction of hydroxylamine reductase (EC 1.7.99.1) since this is a well-known activity of cytochrome cd1.
Synonyms	cd-cytochrome nitrite reductase; [nitrite reductase (cytochrome)] [misleading, see comments.]; cytochrome c-551:O2, NO2+ oxidoreductase; cytochrome cd; cytochrome cd1; hydroxylamine (acceptor) reductase; methyl viologen-nitrite reductase; nitrite reductase (cytochrome; NO-forming)
Product Information	
Form	Liquid or lyophilized powder
EC Number	EC 1.7.2.1
CAS No.	9080-03-9
Reaction	nitric oxide + H2O + ferricytochrome c = nitrite + ferrocytochrome c + 2 H+
Notes	This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

## Storage and Shipping Information

Storage

Store it at +4 °C for short term. For long term storage, store it at -20 °C $\sim$ -80 °C.