

diaminobutyrate acetyltransferase

Cat. No. EXWM-2119

Lot. No. (See product label)

Introduction

Description Requires Na+ or K+ for maximal activity. Ornithine, lysine, aspartate, and α-,β- and

γ-aminobutanoate cannot act as substrates. However, acetyl-CoA can be replaced by propanoyl-CoA, although the reaction proceeds more slowly. Forms part of the ectoine-biosynthesis pathway, the other enzymes involved being EC 2.6.1.76, diaminobutyrate-2-oxoglutarate transaminase and EC 4.2.1.108, ectoine synthase.

Synonyms L-2,4-diaminobutyrate acetyltransferase; L-2,4-diaminobutanoate acetyltransferase;

EctA; diaminobutyric acid acetyltransferase; DABA acetyltransferase; 2,4-

diaminobutanoate acetyltransferase; DAB acetyltransferase; DABAcT; acetyl-CoA:L-

1/1

2,4-diaminobutanoate 4-N-acetyltransferase

Product Information

Form Liquid or lyophilized powder

EC Number EC 2.3.1.178

CAS No. 130456-92-7

Reaction acetyl-CoA + L-2,4-diaminobutanoate = CoA + N4-acetyl-L-2,4-diaminobutanoate

Notes This item requires custom production and lead time is between 5-9 weeks. We can

custom produce according to your specifications.

Storage and Shipping Information

Storage Store it at +4 °C for short term. For long term storage, store it at -20 °C \sim -80 °C.

Tel: 1-631-562-8517 1-516-512-3133 **Email:** info@creative-enzymes.com