

## RING-type E3 ubiquitin transferase

Cat. No. EXWM-2301

Lot. No. (See product label)

### Introduction

**Description** The RING domain of E3 ubiquitin transferase serves as a mediator bringing the ubiquitin-charged E2 ubiquitin-conjugating enzyme and the acceptor protein together to enable the direct transfer of ubiquitin through the formation of an isopeptide bond between the C-terminal glycine residue of ubiquitin and the ε-amino group of an L-lysine residue of the acceptor protein. The RING-E3 domain does not form a catalytic thioester intermediate with ubiquitin (unlike the HECT domain, EC 2.3.2.26). RING-type ubiquitin transferases may occur as single-chain enzymes but also in dimeric forms or in multi-subunit assemblies.

**Synonyms** RING E3 ligase (misleading); ubiquitin transferase RING E3

### Product Information

**Form** Liquid or lyophilized powder

**EC Number** EC 2.3.2.27

**Reaction** S-ubiquitinyl-[E2 ubiquitin-conjugating enzyme]-L-cysteine + [acceptor protein]-L-lysine = [E2 ubiquitin-conjugating enzyme]-L-cysteine + N6-ubiquitinyl-[acceptor protein]-L-lysine

**Notes** This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

### Storage and Shipping Information

**Storage** Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.