

peptidoglycan glycosyltransferase

Cat. No. EXWM-2354

Lot. No. (See product label)

Introduction

- **Description** The enzyme also works when the lysine residue is replaced by meso-2,6-diaminoheptanedioate (meso-2,6-diaminopimelate, A2pm) combined with adjacent residues through its L-centre, as it is in Gram-negative and some Gram-positive organisms. The undecaprenol involved is ditrans, octacis-undecaprenol (for definitions, click here). Involved in the synthesis of cell-wall peptidoglycan.
- SynonymsPG-II; bactoprenyldiphospho-N-acetylmuramoyl-(N-acetyl-D-glucosaminyl)-pentapeptide:peptidoglycan N-
acetylmuramoyl-N-acetyl-D-glucosaminyltransferase; penicillin binding protein (3 or 1B); peptidoglycan
transglycosylase; undecaprenyldiphospho-(N-acetyl-D-glucosaminyl-(1→4)-N-acetyl-D-
muramoylpentapeptide):undecaprenyldiphospho-(N-acetyl-D-glucosaminyl-(1→4)-N-acetyl-D-
muramoylpentapeptide) disaccharidetransferase

Product Information

Form	Liquid or lyophilized powder
EC Number	EC 2.4.1.129
CAS No.	79079-04-2
Reaction	[GlcNAc-(1→4)-Mur2Ac(oyl-L-Ala-γ-D-Glu-L-Lys-D-Ala-D-Ala)]n-diphosphoundecaprenol + GlcNAc-(1→4)- Mur2Ac(oyl-L-Ala-γ-D-Glu-L-Lys-D-Ala-D-Ala)-diphosphoundecaprenol = [GlcNAc-(1→4)-Mur2Ac(oyl-L-Ala-γ- D-Glu-L-Lys-D-Ala-D-Ala)]n+1-diphosphoundecaprenol + undecaprenyl diphosphate
Notes	This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.