Creative Enzymes®

GDP-Man:Man3GlcNAc2-PP-dolichol α-1,2-mannosyltransferase

Cat. No. EXWM-2356

Lot. No. (See product label)

Introduction

Description The biosynthesis of asparagine-linked glycoproteins (N-linked protein glycosylation) utilizes a dolichyl diphosphate-linked glycosyl donor, which is assembled by the series of membrane-bound glycosyltransferases that comprise the dolichol pathway. ALG11 mannosyltransferase from Saccharomyces cerevisiae carries out two sequential steps in the formation of the lipid-linked core oligosaccharide, adding two mannose residues in α(1→2) linkages to the nascent oligosaccharide.

Synonyms ALG11; ALG11 mannosyltransferase; LEW3 (gene name); At2G40190 (gene name); gmd3 (gene name); galactomannan deficiency protein 3; GDP-mannose:glycolipid 1,2-α-D-mannosyltransferase; glycolipid 2-α-mannosyltransferase; GDP-mannose:glycolipid 2-α-D-mannosyltransferase; GDP-Man:Man3GlcNAc2-PP-Dol α-1,2-mannosyltransferase; GDP-α-D-mannose:D-Man-α-(1→3)-[D-Man-α-(1→6)]-D-Man-β-(1→4)-D-GlcNAc-β-(1→4)-D-GlcNAc-diphosphodolichol 2-α-D-mannosyltransferase

Product Information

Form Liquid or lyophilized powder

EC Number EC 2.4.1.131

CAS No. 74506-43-7

Reaction 2 GDP-α-D-mannose + α-D-Man-(1→3)-[α-D-Man-(1→6)]-β-D-Man-(1→4)-β-D-GlcNAc-(1→4)-α-D-GlcNAc-diphosphodolichol = 2 GDP + α-D-Man-(1→2)-α-D-Man-(1→2)-α-D-Man-(1→3)-[α-D-Man-(1→6)]-β-D-Man-(1→4)-β-D-GlcNAc-(1→4)-α-D-GlcNAc-diphosphodolichol

Notes This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage Store it at +4 ℃ for short term. For long term storage, store it at -20 ℃~-80 ℃.