

GDP-Man:Man1GlcNAc2-PP-dolichol α -1,3-mannosyltransferase

Cat. No. EXWM-2357

Lot. No. (See product label)

Introduction

Description The biosynthesis of asparagine-linked glycoproteins utilizes a dolichyl diphosphate-linked glycosyl donor, which is assembled by the series of membrane-bound glycosyltransferases that comprise the dolichol pathway. Alg2 mannosyltransferase from *Saccharomyces cerevisiae* carries out an α 1,3-mannosylation of D-Man- β -(1 \rightarrow 4)-D-GlcNAc- β -(1 \rightarrow 4)-D-GlcNAc-diphosphodolichol, followed by an α 1,6-mannosylation (cf. EC 2.4.1.257), to form the first branched pentasaccharide intermediate of the dolichol pathway.

Synonyms Alg2 mannosyltransferase (ambiguous); ALG2 (gene name, ambiguous); glycolipid 3- α -mannosyltransferase; GDP-mannose:glycolipid 3- α -D-mannosyltransferase; GDP-Man:Man1GlcNAc2-PP-Dol α -1,3-mannosyltransferase; GDP-D-mannose:D-Man- β -(1 \rightarrow 4)-D-GlcNAc- β -(1 \rightarrow 4)-D-GlcNAc-diphosphodolichol 3- α -mannosyltransferase

Product Information

Form Liquid or lyophilized powder

EC Number EC 2.4.1.132

CAS No. 81181-76-2

Reaction GDP- α -D-mannose + β -D-Man-(1 \rightarrow 4)- β -D-GlcNAc-(1 \rightarrow 4)- α -D-GlcNAc-diphosphodolichol = GDP + α -D-Man-(1 \rightarrow 3)- β -D-Man-(1 \rightarrow 4)- β -D-GlcNAc-(1 \rightarrow 4)- α -D-GlcNAc-diphosphodolichol

Notes This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.