

Native Bovine L-Glutamic Dehydrogenase

Cat. No. NATE-0392

Lot. No. (See product label)

Introduction

Description

L-glutamic dehydrogenase catalyzes the conversion of glutamate to α -ketoglutarate. Mammalian forms of this enzyme, including this bovine form, can use either NADP (H) or NAD (H) as coenzymes. L-glutamic dehydrogenase plays a unique role in mammalian metabolism. The reverse reaction catalyzed by this enzyme is the only pathway by which ammonia can become bound to the α -carbon atom of an α -carboxylic acid and thus, is the only source of de novo amino acid synthesis in mammalian species.

Synonyms

glutamic dehydrogenase; glutamate dehydrogenase [NAD (P)]; 9029-12-3; glutamate dehydrogenase [NAD (P)+]; EC 1.4.1.3; L-GLDH; L-Glutamate:NAD[P]+ Oxidoreductase (deaminating)

Product Information

Species

Bovine

Source

Bovine liver

Form

Type I, ammonium sulfate suspension, Suspension in 2.0 M (NH₄)₂SO₄ solution; Type II, lyophilized powder, Contains primarily Citrate buffer salt; Type III, aqueous glycerol solution, Solution in 50% glycerol, pH 7.3.

EC Number

EC 1.4.1.3

CAS No.

9029-12-3

Activity

Type I, > 40 units/mg protein; Type II, > 20 units/mg protein; Type III, > 35 units/mg protein.

Pathway

Alanine, aspartate and glutamate metabolism, organism-specific biosystem; Alanine, aspartate and glutamate metabolism, conserved biosystem; D-Glutamine and D-glutamate metabolism, organism-specific biosystem

Function

ATP binding; GTP binding; glutamate dehydrogenase (NAD⁺) activity

Unit Definition

One unit will reduce 1.0 μ mole of α -ketoglutarate to L-glutamate per min at pH 7.3 at 25°C, in the presence of ammonium ions.

Storage and Shipping Information

Storage

2-8°C