

$\alpha\text{-}1,6\text{-}mannosyl\text{-}glycoprotein }6\text{-}\beta\text{-}N\text{-}acetylglucosaminyltransferase}$

Cat. No. EXWM-2380

Lot. No. (See product label)

Introduction

Description R represents the remainder of the N-linked oligosaccharide in the glycoprotein acceptor (click here for

diagram).

Synonyms N-acetylglucosaminyltransferase V; α -mannoside β -1,6-N-acetylglucosaminyltransferase; uridine

diphosphoacetylglucosamine- α -mannoside $\beta 1 \rightarrow 6$ -acetylglucosaminyltransferase; UDP-N-

acetylglucosamine: α -mannoside- β 1,6 N-acetylglucosaminyltransferase; α -1,3(6)-mannosylglycoprotein β -

1,6-N-acetylglucosaminyltransferase; GnTV

Product Information

Form Liquid or lyophilized powder

EC Number EC 2.4.1.155

CAS No. 83588-90-3

 $\textbf{Reaction} \qquad \text{UDP-N-acetyl-D-glucosamine} + 6-(2-[\text{N-acetyl-}\beta-\text{D-glucosaminyl}] - \alpha-\text{D-mannosyl}) - \beta-\text{D-mannosyl-R} = \text{UDP} + (1-\beta-\text{D-glucosaminyl}) - \beta-\text{D-mannosyl-R} = (1-\beta-\text{D-glucosaminyl}) - \beta-\text{D-glucosaminyl} = (1-\beta-\text{D-glucosaminyl}) - \beta-\text{D-glucosaminyl} = (1-\beta-\text{D-glucosaminyl}) - \beta-\text{D-glucosaminyl} = (1-\beta-\text{D-glucosam$

 $6-(2,6-bis[N-acetyl-\beta-D-glucosaminyl]-\alpha-D-mannosyl)-\beta-D-mannosyl-R$

Notes This item requires custom production and lead time is between 5-9 weeks. We can custom produce

according to your specifications.

Storage and Shipping Information

Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.

Tel: 1-631-562-8517 1-516-512-3133 **Email:** info@creative-enzymes.com

1/1