

glucuronosyl-N-acetylgalactosaminyl-proteoglycan 4- β -N-acetylgalactosaminyltransferase

Cat. No. EXWM-2400

Lot. No. (See product label)

Introduction

Description Involved in the biosynthesis of chondroitin sulfate. The human form of this enzyme

is a bifunctional glycosyltransferase, which also has the 3-β-

glucuronosyltransferase (EC 2.4.1.226, N-acetylgalactosaminyl-proteoglycan 3- β -glucuronosyltransferase) activity required for the synthesis of the chondroitin sulfate disaccharide repeats. Similar chondroitin synthase 'co-polymerases' can be

found in Pasteurella multocida and Escherichia coli.

Synonyms N-acetylgalactosaminyltransferase II; UDP-N-acetyl-D-galactosamine:D-glucuronyl-

N-acetyl-1,3-β-D-galactosaminylproteoglycan β-1,4-N-

acetylgalactosaminyltransferase; chondroitin synthase; glucuronyl-N-

acetylgalactosaminylproteoglycan β-1,4-N-acetylgalactosaminyltransferase; uridine diphosphoacetylgalactosamine-chondroitin acetylgalactosaminyltransferase II;

UDP-N-acetyl-D-galactosamine:β-D-glucuronosyl-($1\rightarrow 3$)-N-acetyl-β-D-galactosaminyl-proteoglycan 4-β-N-acetylgalactosaminyltransferase

Product Information

Form Liquid or lyophilized powder

EC Number EC 2.4.1.175

CAS No. 96189-40-1

Reaction UDP-N-acetyl- α -D-galactosamine + β -D-glucuronosyl- $(1\rightarrow 3)$ -N-acetyl- β -D-

galactosaminyl-proteoglycan = UDP + N-acetyl- β -D-galactosaminyl- $(1\rightarrow 4)$ - β -D-

1/1

 $glucuronosyl\text{-}(1\rightarrow 3)\text{-}N\text{-}acetyl\text{-}\beta\text{-}D\text{-}galactosaminyl\text{-}proteoglycan}$

Notes This item requires custom production and lead time is between 5-9 weeks. We can

custom produce according to your specifications.

Storage and Shipping Information

Store it at +4 °C for short term. For long term storage, store it at -20 °C \sim -80 °C.