

## dolichyl-P-Man:Man6GlcNAc2-PP-dolichol $\alpha$ -1,2-mannosyltransferase

Cat. No. EXWM-2489

Lot. No. (See product label)

## Introduction

**Description** The formation of N-glycosidic linkages of glycoproteins involves the ordered

assembly of the common Glc3Man9GlcNAc2 core-oligosaccharide on the lipid carrier dolichyl diphosphate. Early mannosylation steps occur on the cytoplasmic side of the endoplasmic reticulum with GDP-Man as donor, the final reactions from Man5GlcNAc2-PP-Dol to Man9Glc-NAc2-PP-Dol on the lumenal side use dolichyl  $\beta$ -D-mannosyl phosphate. ALG9 mannosyltransferase catalyses the addition of two different  $\alpha$ -1,2-mannose residues - the addition of  $\alpha$ -1,2-mannose to Man6GlcNAc2-PP-Dol (EC 2.4.1.259) and the addition of  $\alpha$ -1,2-mannose to Man8GlcNAc2-PP-Dol

(EC 2.4.1.261).

Synonyms ALG9; ALG9 α1,2 mannosyltransferase; dolichylphosphomannose-dependent ALG9

mannosyltransferase; ALG9 mannosyltransferase; Dol-P-Man:Man6GlcNAc2-PP-Dol  $\alpha$ -1,2-mannosyltransferase; dolichyl  $\beta$ -D-mannosyl phosphate:D-Man- $\alpha$ -(1 $\rightarrow$ 2)-D-Man- $\alpha$ -(1 $\rightarrow$ 3)-[D-Man- $\alpha$ -(1 $\rightarrow$ 3)-D-Man- $\alpha$ -(1 $\rightarrow$ 4)-D-

GlcNAc- $\beta$ -(1 $\rightarrow$ 4)-D-GlcNAc-diphosphodolichol  $\alpha$ -1,2-mannosyltransferase

## **Product Information**

**Form** Liquid or lyophilized powder

**EC Number** EC 2.4.1.259

**Reaction** dolichyl β-D-mannosyl phosphate + α-D-Man-(1→2)-α-D-Man-(1→2)-α-D-Man-(1→3)-

 $\begin{array}{l} [\alpha\text{-D-Man-}(1\rightarrow 3)-\alpha\text{-D-Man-}(1\rightarrow 6)]-\beta\text{-D-Man-}(1\rightarrow 4)-\beta\text{-D-GlcNAc-}(1\rightarrow 4)-\alpha\text{-D-GlcNAc-} \\ \text{diphosphodolichol} = \alpha\text{-D-Man-}(1\rightarrow 2)-\alpha\text{-D-Man-}(1\rightarrow 2)-\alpha\text{-D-Man-}(1\rightarrow 3)-[\alpha\text{-D-Man-}(1\rightarrow 4)-\beta\text{-D-GlcNAc-}(1\rightarrow 4)-\alpha\text{-D-Man-}(1\rightarrow 4)-\alpha\text{-D-Man$ 

1/1

GlcNAc-diphosphodolichol + dolichyl phosphate

**Notes** This item requires custom production and lead time is between 5-9 weeks. We can

custom produce according to your specifications.

## Storage and Shipping Information

Store it at +4 °C for short term. For long term storage, store it at -20 °C∼-80 °C.