

galactoside 2-α-L-fucosyltransferase

Cat. No. EXWM-2612

Lot. No. (See product label)

Introduction

Description Free lactose can act as acceptor. Normally acts on a glycoconjugate where R (see reaction) is a

glycoprotein or glycolipid. The action on glycolipid was previously listed as EC 2.4.1.89.

Synonyms blood group H α -2-fucosyltransferase; guanosine diphosphofucose-galactoside 2-L-fucosyltransferase; α -

 $(1\rightarrow 2)$ -L-fucosyltransferase; α -2-fucosyltransferase; α -2-L-fucosyltransferase; blood-group substance H-dependent fucosyltransferase; guanosine diphosphofucose-glycoprotein 2- α -fucosyltransferase; guanosine diphosphofucose-lactose fucosyltransferase; GDP fucose-lactose fucosyltransferase; guanosine diphosphofucose-lactose fucosyltransferase; guanosine diphosphofucose- β -D-galactosyl- α -2-L-fucosyltransferase;

guanosine diphosphofucose-galactosylacetylglucosaminylgalactosylglucosylceramide $\alpha\text{-L-}$

fucosyltransferase; guanosine diphosphofucose-glycoprotein $2 - \alpha - L$ -fucosyltransferase; H-gene-encoded β -galactoside $\alpha 1 \rightarrow 2$ fucosyltransferase; secretor-type β -galactoside $\alpha 1 \rightarrow 2$ fucosyltransferase; GDP-L-fucose:lactose fucosyltransferase; GDP- β -L-fucose: β -D-galactosyl-R $2 - \alpha - L$ -

fucosyltransferase

Product Information

Form Liquid or lyophilized powder

EC Number EC 2.4.1.69

CAS No. 56093-23-3

 $\textbf{Reaction} \qquad \text{GDP-}\beta\text{-L-fucose} + \beta\text{-D-galactosyl-}(1→3)\text{-N-acetyl-}\beta\text{-D-glucosaminyl-}(1→3)-\beta\text{-D-galactosyl-}(1→4)$

(1→3)- β -D-galactosyl-(1→4)- β -D-glucosyl-(1↔1)-ceramide

Notes This item requires custom production and lead time is between 5-9 weeks. We can custom produce

according to your specifications.

Storage and Shipping Information

Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.

 1/1