

(Kdo)-lipid IVA 3-deoxy-D-manno-octulosonic acid transferase

Cat. No. EXWM-2697

Lot. No. (See product label)

Introduction

Description

The bifunctional enzyme from Escherichia coli transfers two 3-deoxy-D-manno-oct-2-ulose residues to lipid IVA (cf. EC 2.4.99.12 [lipid IVA 3-deoxy-D-manno-octulosonic acid transferase]). The enzymes from Chlamydia transfer three or more 3-deoxy-D-manno-oct-2-ulose residues and generate genus-specific epitopes.

Synonyms

Kdo transferase; waaA (gene name); kdtA (gene name); 3-deoxy-D-manno-oct-2-ulose acid transferase; 3-deoxy-manno-octulosonic acid transferase; (KDO)-lipid IVA 3-deoxy-D-manno-octulosonic acid transferase

Product Information

Form

Liquid or lyophilized powder

EC Number

EC 2.4.99.13

Reaction

α -Kdo-(2→6)-lipid IVA + CMP-β-Kdo = α -Kdo-(2→4)- α -Kdo-(2→6)-lipid IVA + CMP

Notes

This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage

Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.