

Lysozyme from Human, Recombinant

Cat. No. NATE-0434

Lot. No. (See product label)

Introduction

Description

Lysozymes, also known as muramidase or N-acetylmuramide glycanhydrolase, are glycoside hydrolases. These are enzymes (EC 3.2.1.17) that damage bacterial cell walls by catalyzing hydrolysis of 1,4-beta-linkages between N-acetylmuramic acid and N-acetyl-D-glucosamine residues in a peptidoglycan and between N-acetyl-D-glucosamine residues in chitodextrins. Lysozyme is abundant in a number of secretions, such as tears, saliva, human milk, and mucus. It is also present in cytoplasmic granules of the macrophages and the polymorphonuclear neutrophils (PMNs). Large amounts of lysozyme can be found in egg white. C-type lysozymes are closely related to alpha-lactalbumin in sequence and structure, making them part of the same family. In humans, the lysozyme enzyme is encoded by the LYZ gene.

Synonyms

muramidase; globulin G; mucopeptide glucohydrolase; globulin G1; N,O-diacetylmuramidase; lysozyme g; L-7001; 1,4-N-acetylmuramidase; mucopeptide N-acetylmuramoylhydrolase; PR1-lysozyme; lysozyme; LYZ; LZM; EC 3.2.1.17; 9001-63-2

Product Information

Species

Human

Source

Rice

Form

lyophilized powder

EC Number

EC 3.2.1.17

CAS No.

9001-63-2

Activity

> 100 ,000 units/mg protein (E1%/280)

Pathway

Amyloids, organism-specific biosystem; C-MYB transcription factor network, organism-specific biosystem; Disease, organism-specific biosystem; Salivary secretion, organism-specific biosystem; Salivary secretion, conserved biosystem

Function

hydrolase activity, acting on glycosyl bonds; lysozyme activity

Unit Definition

One unit will produce a ΔA_{450} of 0.001 per min at pH 6.24 at 25°C, using a suspension of *Micrococcus lysodeikticus* as substrate, in a 2.6 mL reaction mixture (1 cm light path).

Usage and Packaging

Package

Package size based on protein content

Storage and Shipping Information

Storage

-70°C