Creative Enzymes®

polymannosyl GlcNAc-diphospho-ditrans,octacis-undecaprenol kinase

Cat. No. EXWM-3013

Lot. No. (See product label)

Introduction

Description	The enzyme is involved in the biosynthesis of the polymannose O-polysaccharide in the outer leaflet of the membrane of Escherichia coli serotype O9a. O-Polysaccharide structures vary extensively because of differences in the number and type of sugars in the repeat unit. The dual kinase/methylase WbdD also catalyses the methylation of 3-phospho-α-D-Man-(1→2)-α-D-Man-(1→2)-α-D-Man-(1→3)-α-D-Man-(1→3)-[α-D-Man-(1→2)-α-D-Man-(1→2)-α-D-Man-(1→3)-α-D-Man-(1→3)]n-α-D-Man-(1→3)-α-D-Man-(1→3)-α-D-GlcNAc-α-diphospho-ditrans,octacis-undecaprenol (cf. EC 2.1.1.294, 3-phospho-α-D-Man-(1→2)-α-D-Man-(1→3)-α-D-Man-(1→3)-α-D-Man-diphospho-ditrans,octacis-undecaprenol 3-phospho-methyltransferase)
Synonyms	WbdD

Product Information

Form	Liquid or lyophilized powder
EC Number	EC 2.7.1.181
Reaction	ATP + α-D-Man-(1→2)-α-D-Man-(1→2)-α-D-Man-(1→3)-α-D-Man-(1→3)-[α-D-Man-(1→2)-α-D-Man-(1→2)-α-D-Man-(1→3)-α-D-Man-(1→3)]n-α-D-Man-(1→3)-α-D-Man-(1→3)-α-D-GlcNAc-diphospho-ditrans,octacis-undecaprenol = ADP + 3-O-phospho-α-D-Man-(1→2)-α-D-Man-(1→2)-α-D-Man-(1→3)-α-D-Man-(1→3)-[α-D-Man-(1→2)-α-D-Man-(1→2)-α-D-Man-(1→3)-α-D-Man-(1→3)]n-α-D-Man-(1→3)-α-D-Man-(1→3)-α-D-GlcNAc-diphospho-ditrans,octacis-undecaprenol
Notes	This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage	Store it at +4 ℃ for short term. For long term storage, store it at -20 ℃~-80 ℃.