

## [hydroxymethylglutaryl-CoA reductase (NADPH)] kinase

Cat. No. EXWM-3152

Lot. No. (See product label)

### Introduction

#### Description

The enzyme is activated by AMP. EC 1.1.1.34, hydroxymethylglutaryl-CoA reductase (NADPH) is inactivated by the phosphorylation of the enzyme protein. Histones can also act as acceptors. The enzyme can also phosphorylate hepatic acetyl-CoA carboxylase (EC 6.4.1.2) and adipose hormone-sensitive lipase (EC 3.1.1.79). Thr-172 within the catalytic subunit ( $\alpha$ -subunit) is the major site phosphorylated by the AMP-activated protein kinase kinase. GTP can act instead of ATP

#### Synonyms

AMPK; AMP-activated protein kinase; HMG-CoA reductase kinase;  $\beta$ -hydroxy- $\beta$ -methylglutaryl-CoA reductase kinase; [hydroxymethylglutaryl-CoA reductase (NADPH<sub>2</sub>)] kinase; 3-hydroxy-3-methylglutaryl coenzyme A reductase kinase; 3-hydroxy-3-methylglutaryl-CoA reductase kinase; hydroxymethylglutaryl coenzyme A reductase kinase; hydroxymethylglutaryl coenzyme A reductase kinase (phosphorylating); hydroxymethylglutaryl-CoA reductase kinase; reductase kinase; STK29

### Product Information

#### Form

Liquid or lyophilized powder

#### EC Number

EC 2.7.11.31

#### CAS No.

172522-01-9

#### Reaction

ATP + [hydroxymethylglutaryl-CoA reductase (NADPH)] = ADP + [hydroxymethylglutaryl-CoA reductase (NADPH)] phosphate

#### Notes

This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

### Storage and Shipping Information

#### Storage

Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.