

Native Potato Acid Phosphatase

Cat. No. NATE-0083

Lot. No. (See product label)

Introduction

Description Native Potato Acid Phosphatase for research on plant phosphatase activity and enzymatic processes.

Perfect for agricultural and biochemical studies. Creative Enzymes provides high-quality solutions.

Applications Phosphatase acid from potato has been used in a study to develop a method of efficient enzymatic

hydrolysis of polyprenyl pyrophosphates. It has also been used in a study to investigate the kinetics of

the hydrolysis of sodium p-nitrophenylphosphate and other phosphoric acid monoesters.

Synonyms acid phosphatase; 9001-77-8; acid phosphomonoesterase; phosphomonoesterase; glycerophosphatase;

acid monophosphatase; acid phosphohydrolase; acid phosphomonoester hydrolase; uteroferrin; acid nucleoside diphosphate phosphatase; orthophosphoric-monoester phosphohydrolase (acid optimum); EC

3.1.3.2; APase

Product Information

Source Potato

Form Type I, Type IV, ammonium sulfate suspension; Suspension in 1.8 M (NH4)2SO4, 10 mM MgCl2, pH 5.5;

Type II, Type III, lyophilized powder.

EC Number EC 3.1.3.2

CAS No. 9001-77-8

Activity Type I, > 200 units/mg protein (biuret); Type II, 0.5-3.0 unit/mg solid; Type III, 3.0-10.0 units/mg solid;

Type IV, > 10.0 units/mg protein (modified Warburg-Christian).

Unit

One unit will hydrolyze 1.0 µmole of p-nitrophenyl phosphate per min at pH 4.8 at 37°C.

Definition

Storage and Shipping Information

Storage 2-8°C

Tel: 1-631-562-8517 1-516-512-3133 **Email:** info@creative-enzymes.com

1/1