

## **Native Rabbit Phosphorylase a**

Cat. No. NATE-0562

Lot. No. (See product label)

## Introduction

Description Phosphorylase A is the active form of glycogen phosphorylase which converts glycogen and

> orthophosphate (Pi) to glucose 1-phoshate (G-1-P). Phosphorylase A can be inhibited by these compounds: Polychlorinated biphenyls, polychlorinated biphenylols and polybrominated biphenyls. Dimeric phosphorylase b is converted to the more active tetramer, phosphorylase a, by the action of

phosphorylase kinase.

Phosphorylase from rabbit muscle has been used in a study to assess the molecular mechanisms of **Applications** 

oleanolic acid. It has also been used in a study to describe the conversion of phosphorylase B to A,

through a conversion enzyme in the presence of 32P-ATP.

Phosphorylase a; EC 2.4.1.1; 9032-10-4; muscle phosphorylase a and b; amylophosphorylase; **Synonyms** 

> polyphosphorylase; amylopectin phosphorylase; glucan phosphorylase;  $\alpha$ -glucan phosphorylase;  $1,4-\alpha$ glucan phosphorylase; glucosan phosphorylase; granulose phosphorylase; maltodextrin phosphorylase; muscle phosphorylase; myophosphorylase; potato phosphorylase; starch phosphorylase;  $1,4-\alpha$ -D-

glucan:phosphate  $\alpha$ -D-glucosyltransferase; phosphorylase (ambiguous)

## **Product Information**

**Species** Rabbit

Source Rabbit muscle

**Form** Lyophilized powder containing β-glycerophosphate and EDTA

EC Number EC 2.4.1.1

CAS No. 9035-74-9

**Purity** 2× crystallization

**Activity** 20-30 units/mg protein

Definition pH 6.8 at 30°C, measured in a system containing phosphoglucomutase, NADP, and glucose-6-phosphate

One unit will form 1.0 μmole of α-D-glucose 1-phosphate from glycogen and orthophosphate per min at

1/1

dehydrogenase. (One µmolar unit is equivalent to ~45 Cori units.)

## Storage and Shipping Information

-20°C Storage

Unit

Tel: 1-631-562-8517 1-516-512-3133 Email: info@creative-enzymes.com