

β-porphyranase

Cat. No. EXWM-3861

Lot. No. (See product label)

Introduction

Description

The backbone of porphyran consists largely (~70%) of (1→3)-linked β-D-galactopyranose followed by (1→4)-linked α-L-galactopyranose-6-sulfate [the other 30% are mostly agarobiose repeating units of (1→3)-linked β-D-galactopyranose followed by (1→4)-linked 3,6-anhydro-α-L-galactopyranose]. This enzyme cleaves the (1→4) linkages between β-D-galactopyranose and α-L-galactopyranose-6-sulfate, forming mostly the disaccharide α-L-galactopyranose-6-sulfate-(1→3)-β-D-galactose, although some longer oligosaccharides of even number of residues are also observed. Since the enzyme is inactive on the non-sulfated agarose portion of the porphyran backbone, some agarose fragments are also included in the products. Methylation of the D-galactose prevents its binding at position -1.

Synonyms

porphyranase; PorA; PorB; endo-β-porphyranase

Product Information

Form

Liquid or lyophilized powder

EC Number

EC 3.2.1.178

Reaction

Hydrolysis of β-D-galactopyranose-(1→4)-α-L-galactopyranose-6-sulfate linkages in porphyran

Notes

This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage

Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.