

UDP-N,N'-diacetylbacillosamine 2-epimerase (hydrolysing)

Cat. No. EXWM-3868

Lot. No. (See product label)

Introduction

Description

Requires Mg²⁺. Involved in biosynthesis of legionaminic acid, a nonulosonate derivative that is incorporated by some bacteria into assorted virulence-associated cell surface glycoconjugates. The initial product formed by the enzyme from *Legionella pneumophila*, which incorporates legionaminic acid into the O-antigen moiety of its lipopolysaccharide, is 2,4-diacetamido-2,4,6-trideoxy- α -D-mannopyranose, which rapidly mutarotates to a mixture of anomers. The enzyme from *Campylobacter jejuni*, which incorporates legionaminic acid into flagellin, prefers GDP-N,N'-diacetylbacillosamine.

Synonyms

UDP-Bac2Ac4Ac 2-epimerase; NeuC

Product Information

Form

Liquid or lyophilized powder

EC Number

EC 3.2.1.184

Reaction

UDP-N,N'-diacetylbacillosamine + H₂O = UDP + 2,4-diacetamido-2,4,6-trideoxy-D-mannopyranose

Notes

This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage

Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.