

ginsenosidase type IV

Cat. No. EXWM-3878 Lot. No. (See product label)

| Introduction | |
|----------------------------------|---|
| Description | Ginsenosidase type IV catalyses the sequential hydrolysis of the 6-O- β -D-(1 \rightarrow 2)- glycosidic bond or the 6-O- α -D-(1 \rightarrow 2)-glycosidic bond in protopanaxatriol-type ginsenosides with a disacchride attached to the C6 position, followed by the hydrolysis of the remaining 6-O- β -D-glycosidic bond (e.g. ginsenoside Re \rightarrow ginsenoside Rg1 \rightarrow ginsenoside F1). |
| Product Information | |
| Form | Liquid or lyophilized powder |
| EC Number | EC 3.2.1.194 |
| Reaction | a protopanaxatriol-type ginsenoside with two glycosyl residues at position $6 + 2$ H2O = a protopanaxatriol-type ginsenoside with no glycosidic modification at position $6 + D$ -glucopyranose + a monosaccharide (overall reaction); (1a) a protopanaxatriol-type ginsenoside with two glycosyl residues at position $6 + H2O =$ a protopanaxatriol-type ginsenoside with a single glucosyl at position $6 + a$ monosaccharide; (1b) a protopanaxatriol-type ginsenoside with a single glucosyl at position $6 + H2O =$ a protopanaxatriol-type ginsenoside with no glycosidic modification at position $6 + D$ -glucopyranose |
| Notes | This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications. |
| Storage and Shipping Information | |

Storage and Shipping Information

Storage

Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.