

limonene-1,2-epoxide hydrolase

Cat. No. EXWM-4005

Lot. No. (See product label)

Introduction

Description

Involved in the monoterpene degradation pathway of the actinomycete *Rhodococcus erythropolis*. The enzyme hydrolyses several alicyclic and 1-methyl-substituted epoxides, such as 1-methylcyclohexene oxide, indene oxide and cyclohexene oxide. It differs from the previously described epoxide hydrolases [EC 3.3.2.4 (trans-epoxysuccinate hydrolase), EC 3.3.2.6 (leukotriene-A4 hydrolase), EC 3.3.2.7 (hepoxilin-epoxide hydrolase), EC 3.3.2.9 (microsomal epoxide hydrolase) and EC 3.3.2.10 (soluble epoxide hydrolase)] as it is not inhibited by 2-bromo-4'-nitroacetophenone, diethyl dicarbonate, 4-fluorochalcone oxide or 1,10-phenanthroline. Both enantiomers of menth-8-ene-1,2-diol [i.e. (1R,2R,4S)-menth-8-ene-1,2-diol and (1S,2S,4R)-menth-8-ene-1,2-diol] are metabolized.

Synonyms

limonene oxide hydrolase

Product Information

Form

Liquid or lyophilized powder

EC Number

EC 3.3.2.8

CAS No.

216503-88-7

Reaction

1,2-epoxymenth-8-ene + H₂O = menth-8-ene-1,2-diol

Notes

This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage

Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.