

Xaa-Xaa-Pro tripeptidyl-peptidase

Cat. No. EXWM-4046

Lot. No. (See product label)

Introduction

Description This cell-surface-associated serine exopeptidase is found in the Gram-negative, anaerobic bacterium

Porphyromonas gingivalis, which has been implicated in adult periodontal disease. The enzyme releases the N-terminal tripeptide of peptides, such as interleukin-6. It has an absolute requirement for a proline residue at the P1 position but is completely inactivated by a proline residue at the P1' position. The size of

the peptide does not affect the rate of reaction.

Synonyms prolyltripeptidyl amino peptidase; prolyl tripeptidyl peptidase; prolyltripeptidyl aminopeptidase; PTP-A;

TPP

Product Information

Form Liquid or lyophilized powder

EC Number EC 3.4.14.12

Reaction Hydrolysis of Xaa-Xaa-Pro-Yaa- releasing the N-terminal tripeptide of a peptide with Pro as the third

residue (position P1) and where Yaa is not proline

Notes This item requires custom production and lead time is between 5-9 weeks. We can custom produce

according to your specifications.

Storage and Shipping Information

Store it at +4 °C for short term. For long term storage, store it at -20 °C∼-80 °C.

Tel: 1-631-562-8517 1-516-512-3133 **Email:** info@creative-enzymes.com

1/1