

glutamate carboxypeptidase II

Cat. No. EXWM-4074

Lot. No. (See product label)

Introduction

Description A metallo-carboxypeptidase that is predominantly expressed as a membrane-bound enzyme of 94-100 kDa, but also exists in a soluble form. Hydrolyses α -peptide bonds in Ac-Asp-Glu, Asp-Glu, and Glu-Glu, but also γ -glutamyl bonds in γ -Glu-Glu, and folylpoly- γ -glutamates. With folylpoly- γ -glutamates, shows processive carboxypeptidase activity to produce pteroylmonoglutamate. Does not hydrolyse Ac- β -Asp-Glu. Known inhibitors: quisqualic acid, Ac- β -Asp-Glu, and 2-phosphonomethyl-pentanedioate. In peptidase family M28 of *Vibrio leucyl* aminopeptidase. The release of C-terminal glutamate from folylpoly- γ -glutamates is also catalysed by EC 3.4.17.11 (glutamate carboxypeptidase) and EC 3.4.19.9 (γ -Glu-X carboxypeptidase).

Synonyms N-acetylated- γ -linked-acidic dipeptidase (NAALADase); folate hydrolase; prostate-specific membrane antigen; pteroylpoly- γ -glutamate carboxypeptidase; microsomal γ -glutamyl carboxypeptidase; pteroylpolyglutamate hydrolase; folylpolyglutamate hydrolase; pteroylpoly- γ -glutamate hydrolase; pteroylpolyglutamyl hydrolase; pteroylpolyglutamate hydrolase; pteroylpolyglutamic acid hydrolase; PSM antigen; acetylaspartylglutamate dipeptidase; NAALA dipeptidase; rat NAAG peptidase; mGCP; membrane glutamate carboxypeptidase; N-acetylated- α -linked-amino dipeptidase; prostrate-specific membrane antigen; N-Acetylated α -linked acidic dipeptidase; PSMA

Product Information

Form Liquid or lyophilized powder

EC Number EC 3.4.17.21

CAS No. 9074-87-7

Reaction Release of an unsubstituted, C-terminal glutamyl residue, typically from Ac-Asp-Glu or folylpoly- γ -glutamates

Notes This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.