

ubiquitinyl hydrolase 1

Cat. No. EXWM-4084

Lot. No. (See product label)

Introduction

Description Links to polypeptides smaller than 60 residues are hydrolysed more readily than those to larger polypeptides. Isoforms exist with quantitatively different specificities, amongst the best known being UCH-L1 and UCH-L3, which are major proteins of the brain of mammals. Inhibited by ubiquitin aldehyde (in which Gly76 is replaced by aminoacetaldehyde). Ubiquitinyl hydrolase 1 is the type example of peptidase family C12, with a similar protein fold to papain and catalytic amino acids Cys, His and Asp. There is a separate family (C19) of enzymes that also hydrolyse ubiquitinyl bonds, and it is thought that all the ubiquitinyl hydrolases are also ubiquitin thiolesterases (EC 3.1.2.15)

Synonyms ubiquitin C-terminal hydrolase; yeast ubiquitin hydrolase

Product Information

Form Liquid or lyophilized powder

EC Number EC 3.4.19.12

CAS No. 86480-67-3, 189642-63-5

Reaction Thiol-dependent hydrolysis of ester, thioester, amide, peptide and isopeptide bonds formed by the C-terminal Gly of ubiquitin (a 76-residue protein attached to proteins as an intracellular targeting signal)

Notes This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.