

## N-acetylglucosaminylphosphatidylinositol deacetylase

Cat. No. EXWM-4480

Lot. No. (See product label)

## Introduction

**Description** Involved in the second step of glycosylphosphatidylinositol (GPI) anchor formation in all eukaryotes. The

enzyme appears to be composed of a single subunit (PIG-L in mammalian cells and GPI12 in yeast). In some species, the long-chain sn-1-acyl group of the phosphatidyl group is replaced by a long-chain alkyl

or alk-1-enyl group.

**Synonyms** N-acetyl-D-glucosaminylphosphatidylinositol acetylhydrolase; N-acetylglucosaminylphosphatidylinositol

de-N-acetylase; GlcNAc-PI de-N-acetylase; GlcNAc-PI deacetylase; acetylglucosaminylphosphatidylinositol

deacetylase

## **Product Information**

**Form** Liquid or lyophilized powder

**EC Number** EC 3.5.1.89

**CAS No.** 122191-30-4

**Reaction**  $6-(N-acetyl-\alpha-D-glucosaminyl)-1-phosphatidyl-1D-myo-inositol + H2O = <math>6-(\alpha-D-glucosaminyl)-1-phosphatidyl-1D-myo-inositol + H2O = (6-(\alpha-D-glucosaminyl)-1-phosphatidyl-1D-myo-inositol + H2O = (6-(\alpha-D-glucosaminyl)-1-phosphatidyl-1D-myo-inositol + H2O = (6-(\alpha-D-glucosaminyl)-1-phosphatidyl-1D-myo-inositol + H2O = (6-(\alpha-D-glucosaminyl)-1-phosphatidyl-1D-myo-inositol + H2O = (6-(\alpha-D-gluc$ 

phosphatidyl-1D-myo-inositol + acetate

**Notes** This item requires custom production and lead time is between 5-9 weeks. We can custom produce

according to your specifications.

## Storage and Shipping Information

Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.

**Tel:** 1-631-562-8517 1-516-512-3133 **Email:** info@creative-enzymes.com

1/1