

1,5-anhydro-D-fructose dehydratase

Cat. No. EXWM-4951

Lot. No. (See product label)

Introduction

Description

This enzyme catalyses one of the steps in the anhydrofructose pathway, which leads to the degradation of glycogen and starch via 1,5-anhydro-D-fructose. The other enzymes involved in this pathway are EC 4.2.1.110 (aldos-2-ulose dehydratase), EC 4.2.2.13 [exo-(1→4)-α-D-glucan lyase] and EC 5.3.2.7 (ascopyrone tautomerase). Requires divalent (Ca²⁺ or Mg²⁺) or monovalent cations (Na⁺) for optimal activity. Unlike EC 4.2.1.110, the enzyme is specific for 1,5-anhydro-D-fructose as substrate and shows no activity towards aldose-2-uloses such as 2-dehydroglucose. In addition, it is inhibited by its end-product ascopyrone M and it cannot convert ascopyrone M into microthecin, as can EC 4.2.1.110.

Synonyms

1,5-anhydro-D-fructose 4-dehydratase; 1,5-anhydro-D-fructose hydrolyase; 1,5-anhydro-D-arabino-hex-2-ulose dehydratase; AFDH; AF dehydratase; 1,5-anhydro-D-fructose hydro-lyase

Product Information

Form

Liquid or lyophilized powder

EC Number

EC 4.2.1.111

Reaction

1,5-anhydro-D-fructose = 1,5-anhydro-4-deoxy-D-glycero-hex-3-en-2-ulose + H₂O

Notes

This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage

Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.