

UDP-N-acetylglucosamine 4,6-dehydratase (configurationinverting)

Cat. No. EXWM-4955 Lot. No. (See product label)

Introduction	
Description	Contains NADP+ as a cofactor. This is the first enzyme in the biosynthetic pathway of pseudaminic acid, a sialic-acid-like sugar that is unique to bacteria and is used by Helicobacter pylori to modify its flagellin. This enzyme plays a critical role in H. pylori's pathogenesis, being involved in the synthesis of both functional flagella and lipopolysaccharides. It is completely inhibited by UDP- α -D-galactose. The reaction results in the chirality of the C-5 atom being inverted. It is thought that Lys-133 acts sequentially as a catalytic acid, protonating the C-6 hydroxy group and as a catalytic base, abstracting the C-5 proton, resulting in the elimination of water. This enzyme belongs to the short-chain dehydrogenase/reductase family of enzymes.
Synonyms	FlaA1; UDP-N-acetylglucosamine 5-inverting 4,6-dehydratase; PseB; UDP-N- acetylglucosamine hydro-lyase (inverting; UDP-2-acetamido-2,6-dideoxy-β-L- arabino-hex-4-ulose-forming)
Product Information	
Form	Liquid or lyophilized powder
EC Number	EC 4.2.1.115
Reaction	UDP-N-acetyl- α -D-glucosamine = UDP-2-acetamido-2,6-dideoxy- β -L-arabino-hex-4-ulose + H2O
Notes	This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage

Store it at +4 ℃ for short term. For long term storage, store it at -20 ℃~-80 ℃.