

## hydroperoxy icosatetraenoate dehydratase

Cat. No. EXWM-4993

Lot. No. (See product label)

### Introduction

#### Description

Binds Fe<sup>2+</sup>. The mammalian enzymes accept a range of hydroperoxyicosatetraenoates (HPETE). The human enzyme has highest activity with (12R)-HPETE, followed by (12S)-HPETE and (15R)-HPETE with much lower efficiency. The murine enzyme has highest activity with (8R)-HPETE followed by (8S)-HPETE. All HPETE isoforms are converted to the corresponding oxoicosatetraenoate forms (KETE). The enzymes also catalyse the reaction of EC 5.4.4.7, hydroperoxy icosatetraenoate isomerase.

#### Synonyms

epidermal lipxygenase-3 (ambiguous); eLOX3 (ambiguous)

### Product Information

#### Form

Liquid or lyophilized powder

#### EC Number

EC 4.2.1.152

#### Reaction

a hydroperoxyicosatetraenoate = an oxoicosatetraenoate + H<sub>2</sub>O

#### Notes

This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

### Storage and Shipping Information

#### Storage

Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.