

chondroitin AC lyase

Cat. No. EXWM-5104

Lot. No. (See product label)

Introduction

Description Acts on chondroitin 4-sulfate and chondroitin 6-sulfate, but less well on hyaluronate. In general, chondroitin sulfate (CS) and dermatan sulfate (DS) chains comprise a linkage region, a chain cap and a repeat region. The repeat region of CS is a repeating disaccharide of glucuronic acid (GlcA) and N-acetylgalactosamine (GalNAc) [-4)GlcA(β1-3)GalNAc(β1-)]_n, which may be O-sulfated on the C-4 and/or C-6 of GalNAc and C-2 of GlcA. GlcA residues of CS may be epimerized to iduronic acid (IdoA) forming the repeating disaccharide [-4)IdoA(α1-3)GalNAc(β1-)]_n of DS. Both the concentrations and locations of sulfate-ester substituents vary with glucosaminoglycan source.

Synonyms chondroitinase (ambiguous); chondroitin sulfate lyase; chondroitin AC eliminase; chondroitinase AC; ChnAC

Product Information

Form Liquid or lyophilized powder

EC Number EC 4.2.2.5

CAS No. 9047-57-8

Reaction Eliminative degradation of polysaccharides containing 1,4-β-D-hexosaminy and 1,3-β-D-glucuronosyl linkages to disaccharides containing 4-deoxy-β-D-gluc-4-enuronosyl groups

Notes This item requires custom production and lead time is between 5-9 weeks. We can custom produce according to your specifications.

Storage and Shipping Information

Storage Store it at +4 °C for short term. For long term storage, store it at -20 °C~-80 °C.