

β-Glucanase 1, thermostable, Recombinant

Cat. No. NATE-0764

Lot. No. (See product label)

Introduction

Description Beta-glucosidase is a glucosidase enzyme located in on the brush border of the small intestine that acts upon β1->4 bonds linking two glucose or glucose-substituted molecules (i.e., the disaccharide cellobiose). It is one of the cellulases, enzymes involved in the decomposition of cellulose and related polysaccharides; more specifically, an exocellulase with specificity for a variety of beta-D-glycoside substrates. It catalyzes the hydrolysis of terminal non-reducing residues in beta-D-glucosides with release of glucose.

Applications β-Glucanase is used to study cell wall modifications and for carbohydrate hydrolysis. It has been used to supplement barley-based diets for poultry and to study the reopening signal conduits and release of dormancy in the Populus species.

Synonyms β-Glucanase 1; β-Glucanase 1, thermostable; 62213-14-3

Product Information

Source E. coli

Form liquid, Supplied as a solution in 50 mM Tris-HCl, pH 8.0, 100 mM NaCl, and 25% glycerol.

CAS No. 62213-14-3

Molecular Weight mol wt 45 kDa

Purity > 90% (SDS-PAGE) 19-21 mg protein/mL (280 nm, UV)

Activity > 10 units/mg protein

Unit Definition One unit will produce 1 μmole of reducing sugar (measured as glucose) from Beta-glucan per minute at pH 5.8 at 70°C.

Storage and Shipping Information

Storage 2-8°C