

Native Trichoderma sp. Laminarinase

Cat. No. NATE-0377 Lot. No. (See product label)

| Introduction | |
|---------------------|---|
| Description | β -glucanases degrade β -1,4-glucans of cellulose, xyloglucan and β -1,4-xylan. β - Glucanase represents a group of carbohydrate enzymes which break down glycosidic bonds within beta-glucan. It forms the main constituent of fungal cell walls and could be a potential structural and storage polysaccharide of marine macro-algae. It has the ability to degrade fungal cell walls and may be involved in defense mechanism of plants against pathogenic fungi. |
| Synonyms | endo-1,3-β-D-glucanase; laminarinase; laminaranase; β-1,3-glucanase; β-1,3-1,4- glucanase; endo-1,3-β-glucanase; endo-β-1,3 (4)-glucanase; endo-β-1,3-1,4- glucanase; endo-β-(1 \rightarrow 3)-D-glucanase; endo-1,3-1,4-β-D-glucanase; endo-β-(1-3)-D- glucanase; endo-β-1,3-glucanase IV; endo-1,3-β-D-glucanase; 1,3-(1,3; 1,4)-β-D- glucan 3 (4)-glucanohydrolase; EC 3.2.1.6; 9074-98-0 |
| Product Information | |
| Source | Trichoderma sp. |
| Form | powder |
| EC Number | EC 3.2.1.6 |
| CAS No. | 62213-14-3 |
| Activity | 100-400 units/g solid |
| Unit Definition | One unit will liberate 1.0 mg of reducing sugar (measured as glucose) from laminarin per min at pH 5.0 at 37°C. |
| | |

Storage and Shipping Information

Storage 2-8°C